## An Anisotropic, Non-Singular Early Universe Model

#### Pierre-Philippe Dechant

Astrophysics Group, Cavendish Laboratory, Cambridge University UniverseNet - Poster session

September 22, 2008

Pierre-Philippe Dechant An Anisotropic, Non-Singular Early Universe Model

• Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon

- Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon
- Biaxial Bianchi IX with scalar field matter

- Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon
- Biaxial Bianchi IX with scalar field matter
- Pancaking solution with  $R_1 \propto t$ ,  $R_2 = R_3 \propto \text{const}$ ,  $\phi = \text{const}$ through the 'Big Bang' at t = 0

- Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon
- Biaxial Bianchi IX with scalar field matter
- Pancaking solution with  $R_1 \propto t$ ,  $R_2 = R_3 \propto \text{const}$ ,  $\phi = \text{const}$ through the 'Big Bang' at t = 0
- No longer singular: curvature is bounded, (some) geodesics go through; parity inverted on 'the other side'

- Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon
- Biaxial Bianchi IX with scalar field matter
- Pancaking solution with  $R_1 \propto t$ ,  $R_2 = R_3 \propto \text{const}$ ,  $\phi = \text{const}$ through the 'Big Bang' at t = 0
- No longer singular: curvature is bounded, (some) geodesics go through; parity inverted on 'the other side'
- Model isotropises and inflates.  $n_s$  and r compatible with observations; low- $\ell$  dip; reasonably stable around biaxiality

- Pierre-Philippe Dechant, Anthony N. Lasenby, Michael P. Hobson: on the arxiv soon
- Biaxial Bianchi IX with scalar field matter
- Pancaking solution with  $R_1 \propto t$ ,  $R_2 = R_3 \propto \text{const}$ ,  $\phi = \text{const}$ through the 'Big Bang' at t = 0
- No longer singular: curvature is bounded, (some) geodesics go through; parity inverted on 'the other side'
- Model isotropises and inflates.  $n_s$  and r compatible with observations; low- $\ell$  dip; reasonably stable around biaxiality
- Structure on the largest scales could have formed when the universe was still oblate